

VIRTUAL BETTI NUMBERS OF REAL ALGEBRAIC VARIETIES

CLINT MCCRORY AND ADAM PARUSIŃSKI

ABSTRACT. The weak factorization theorem for birational maps is used to prove that for all $i \geq 0$ the i th mod 2 Betti number of compact nonsingular real algebraic varieties has a unique extension to a *virtual Betti number* β_i defined for all real algebraic varieties, such that if Y is a closed subvariety of X , then $\beta_i(X) = \beta_i(X \setminus Y) + \beta_i(Y)$.

We define an invariant of the Grothendieck ring $K_0(\mathcal{V}_{\mathbb{R}})$ of real algebraic varieties, the *virtual Poincaré polynomial* $\beta(X, t)$, which is a ring homomorphism $K_0(\mathcal{V}_{\mathbb{R}}) \rightarrow \mathbb{Z}[t]$. For X nonsingular and compact, $\beta(X, t)$ is the classical Poincaré polynomial for cohomology with $\mathbb{Z}_2$ coefficients. The coefficients of the virtual Poincaré polynomial are the virtual Betti numbers. By the weak factorization theorem for birational morphisms (Abramovich *et al.* [1]), the existence of $\beta(X, t)$ follows from a simple formula for the Betti numbers of the blowup of a compact nonsingular variety along a closed nonsingular center. The existence of the virtual Betti numbers for certain real analytic spaces, including real algebraic varieties, has also been announced by Totaro [23].

Kontsevich's motivic measure on the arc space of a complex algebraic variety takes values in the completion of the localized Grothendieck ring (*cf.* [20], [11], [7]). Completion with respect to virtual dimension is possible because complex varieties which are equivalent in the Grothendieck ring have the same dimension. Using the virtual Poincaré polynomial we prove that dimension is an invariant of the Grothendieck ring of real algebraic varieties. It follows that motivic measures can be defined on arc spaces of real varieties.

By a real algebraic variety we mean an algebraic variety in the sense of Serre [22] over the field $\mathbb{R}$ of real numbers. For our purposes there is no loss of generality in restricting our attention to affine real algebraic varieties; over $\mathbb{R}$ every quasiprojective variety is affine. For background on real algebraic varieties we refer the reader to [4].

1. GENERALIZED EULER CHARACTERISTICS

Definition 1.1. The *Grothendieck group* of real algebraic varieties is the abelian group generated by symbols $[X]$, where X is a real algebraic variety, with the two relations

- (1) $[X] = [Y]$ if X and Y are isomorphic,
- (2) $[X] = [X \setminus Y] + [Y]$ if Y is a closed subvariety of X .

The product of varieties induces a ring structure

- (3) $[X] \cdot [Y] = [X \times Y]$,

and the resulting ring, denoted by $K_0(\mathcal{V}_{\mathbb{R}})$, is called the *Grothendieck ring* of real algebraic varieties. The class of a point is the unit of $K_0(\mathcal{V}_{\mathbb{R}})$. The zero element of $K_0(\mathcal{V}_{\mathbb{R}})$ is the class of the empty set.

Definition 1.2. A *generalized (proper) Euler characteristic* of real algebraic varieties, with values in a ring R , is a ring homomorphism $e : K_0(\mathcal{V}_{\mathbb{R}}) \rightarrow R$. In other words, for all varieties X and Y ,

- (1) $e(X) = e(Y)$ if X and Y are isomorphic,
- (2) $e(X) = e(X \setminus Y) + e(Y)$ if Y is a closed subvariety of X ,
- (3) $e(X \times Y) = e(X) \cdot e(Y)$.

The Euler characteristic with compact supports χ_c is an example of such a homomorphism. In fact the only generalized Euler characteristics which are homeomorphism invariants are constant multiples of χ_c (cf. [21]). The standard Euler characteristic χ does not satisfy the additivity property 1.2(2) for real algebraic varieties, and hence it is *not* a generalized Euler characteristic.

For example, let X be the unit circle in the plane, and let Y be a point of X . Then $\chi(X) = 0$, but $\chi(X \setminus Y) = 1$ and $\chi(Y) = 1$. On the other hand, $\chi_c(X) = 0$, $\chi_c(X \setminus Y) = -1$ and $\chi_c(Y) = 1$.

There are several ways to view the Euler characteristic with compact supports. By definition $\chi_c(X)$ is the Euler characteristic of the sheaf cohomology of X with compact supports, with coefficients in the constant sheaf $\mathbb{Z}$. (Below we use coefficients $\mathbb{Z}_2$, the integers mod 2.) We denote by $H_c^i(X)$ the i th cohomology of X with compact supports. Classical cohomology has arbitrary closed supports. Property 1.2(2) for χ_c follows from the long exact cohomology sequence of the pair (X, Y) ([15], 4.10). If X is triangulated then the sheaf cohomology of X with compact supports is isomorphic to the simplicial cohomology of X with compact supports.

For coefficients in a field the i th cohomology of X with compact supports is isomorphic to the i th Borel-Moore homology of X ([5], 3.3). Thus $\chi_c(X)$ is also equal to the Euler characteristic of the Borel-Moore homology of X . Borel-Moore homology has arbitrary closed supports. Classical homology has compact supports. If X is triangulated the Borel-Moore homology of X is isomorphic to the simplicial homology of X with closed supports (*i.e.* using possibly infinite simplicial chains). If $\bar{X}$ is a compact space containing X such that $\bar{X}$ is triangulated with $\bar{X} \setminus X$ as a subcomplex, then the Borel-Moore homology of X is isomorphic to the relative simplicial homology of the pair $(\bar{X}, \bar{X} \setminus X)$. Every real algebraic variety X admits a compactification with such a triangulation (cf. [19], [17], [4]).

Let X be a compact nonsingular real algebraic variety and let C be a closed nonsingular subvariety of X . Denote by $\text{Bl}_C X \rightarrow X$ the blow-up of X with center C and by E the exceptional divisor. Then since $\text{Bl}_C X \setminus E$ is isomorphic to $X \setminus C$, we have

$$(1.1) \quad [\text{Bl}_C X] - [E] = [X] - [C].$$

Bittner [3] has proved using the weak factorization theorem of Abramovich *et al.* [1] that one may replace 1.1(2) in the definition of the Grothendieck ring by (1.1). The following result is equivalent to Bittner's, and our proof is a reorganization of hers.

Theorem 1.3. *Let $X \mapsto e(X)$ be a function defined on compact nonsingular real algebraic varieties with values in a abelian group G , such that $e(X) = e(Y)$ if X and Y are isomorphic. Suppose that $e(\emptyset) = 0$ and for any blow-up $\text{Bl}_C X \rightarrow X$, where X is a compact nonsingular real algebraic variety and $C \subset X$ a closed nonsingular subvariety, with exceptional divisor E ,*

$$e(\text{Bl}_C X) - e(E) = e(X) - e(C).$$

Then e extends uniquely to a group homomorphism $K_0(\mathcal{V}_{\mathbb{R}}) \rightarrow G$.

If e takes values in a ring R and $e(X \times Y) = e(X) \cdot e(Y)$ for all X, Y compact nonsingular then this extension of e is a ring homomorphism $K_0(\mathcal{V}_{\mathbb{R}}) \rightarrow R$.

Proof. First we note that if X is the disjoint union of the compact nonsingular varieties Y and Z ,

$$X = Y \sqcup Z,$$

then

$$e(X) = e(Y) + e(Z).$$

For the blow-up of X with center $C = Y$ is Z , and the exceptional divisor E is empty (cf. [4], p. 78 *et seq.*).

We prove the following statement by induction on n :

(F_n) *There is a unique map $X \mapsto \varphi_n(X) \in G$ defined on varieties of dimension $\leq n$ such that*

- (a) $\varphi_n(X) = \varphi_n(Y)$ if X and Y are isomorphic,
- (b) $\varphi_n(X) = e(X)$ for X compact and nonsingular,
- (c) $\varphi_n(X \setminus Y) = \varphi_n(X) - \varphi_n(Y)$ if Y is a closed subvariety of X .

The case $n = 0$ is trivial. Suppose (F_{n-1}) holds. If $\dim X \leq n - 1$ then we put $\varphi_n(X) := \varphi_{n-1}(X)$. If $\dim X = n$ then $\varphi_n(X)$ is defined follows:

- (i) Let X be nonsingular and let $\bar{X}$ be a nonsingular compactification of X . Then

$$\varphi_n(X) := e(\bar{X}) - \varphi_n(\bar{X} \setminus X),$$

- (ii) Let X be singular and let $X = \bigsqcup S_i$ be a stratification of X . Then

$$\varphi_n(X) := \sum \varphi_n(S_i).$$

Here by a nonsingular compactification of the variety X we mean a compact nonsingular variety $\bar{X}$ which contains a subvariety U isomorphic to X such that the Zariski closure of U in $\bar{X}$ equals $\bar{X}$.

By a stratification $\mathcal{S} = \{S_i\}$ of X we mean a finite disjoint union $X = \bigsqcup S_i$ where each stratum S_i is a locally closed nonsingular subvariety of X . Denote by $\overline{S_i}^Z$ the Zariski closure of S_i in X . We also require that, for every stratum S_i , the set $\overline{S_i}^Z \setminus S_i$ is a closed algebraic subvariety of X which is a union of strata of $\mathcal{S}$.

First we show that the definition (i) of $\varphi_n(X)$ for X nonsingular is independent of the choice of compactification. Let $\bar{X}$ and $\bar{X}'$ be two nonsingular compactifications of X . Then $\bar{X}$ and $\bar{X}'$ are birationally isomorphic. By the weak factorization theorem there is a finite sequence of blow-ups and blow-downs with nonsingular centers joining $\bar{X}$ and $\bar{X}'$:

$$\bar{X} \xleftarrow{\varphi_0} X_1 \xrightarrow{\varphi_1} X_2 \longleftarrow \cdots \longrightarrow X_{k-1} \xleftarrow{\varphi_{k-1}} X_k \xrightarrow{\varphi_k} \bar{X}'$$

Thus to prove independence we may suppose that $\bar{X}' = \text{Bl}_C \bar{X}$, where C is nonsingular and $C \cap X = \emptyset$. Let E denote the exceptional divisor. Then

$$\begin{aligned} e(\bar{X}') - \varphi_n(\bar{X}' \setminus X) &= e(\bar{X}') - \varphi_n(E) - \varphi_n(\bar{X}' \setminus X \setminus E) \\ &= e(\bar{X}') - \varphi_n(E) - \varphi_n(\bar{X} \setminus X \setminus C) = e(\bar{X}') - \varphi_n(E) + \varphi_n(C) - \varphi_n(\bar{X} \setminus X) \\ &= (e(\bar{X}') - e(E) + e(C)) - \varphi_n(\bar{X} \setminus X) = e(\bar{X}) - \varphi_n(\bar{X} \setminus X). \end{aligned}$$

Next we show that $\varphi_n(X \setminus Y) = \varphi_n(X) - \varphi_n(Y)$ for X and Y nonsingular. Let $\bar{X}$ be a nonsingular compactification of X . Let $D = \bar{X} \setminus X$. (It is not necessary to assume that D is a divisor in $\bar{X}$.) First consider the case $\overline{X \setminus Y}^Z = X$, i.e. Y contains no irreducible component of X . Since $\dim Y < n$, we have

$$\varphi_n(X \setminus Y) = e(\bar{X}) - \varphi_n(D \cup Y) = e(\bar{X}) - \varphi_n(D) - \varphi_n(Y) = \varphi_n(X) - \varphi_n(Y).$$

The next case is when Y is the union of some irreducible components of X . Let $\bar{Y}$ denote the Zariski closure of Y in $\bar{X}$. Clearly $\bar{Y}$ is a union of irreducible components of $\bar{X}$. Let $W = \bar{X} \setminus \bar{Y}$. Then

$$\begin{aligned} \varphi_n(X \setminus Y) &= e(W) - \varphi_n(D \cap W) \\ &= e(\bar{X}) - e(\bar{Y}) - \varphi_n(D) + \varphi_n(D \cap \bar{Y}) = \varphi_n(X) - \varphi_n(Y). \end{aligned}$$

In the general case we denote by X' the union of the irreducible components of X contained in Y and by X'' the union of the remaining irreducible components. Let $Y'' = Y \cap X''$. Then by the above cases

$$\begin{aligned} \varphi_n(X \setminus Y) &= \varphi_n(X'' \setminus Y'') = \varphi_n(X'') - \varphi_n(Y'') \\ &= \varphi_n(X) - \varphi_n(X') - (\varphi_n(Y) - \varphi_n(X')) = \varphi_n(X) - \varphi_n(Y). \end{aligned}$$

In particular, suppose X is nonsingular, $\dim X \leq n$, and let $X = \bigsqcup S_i$ be a stratification of X . Let S_0 be a stratum of smallest dimension. Then S_0 is a closed nonsingular variety of X and hence

$$\varphi_n(X \setminus S_0) = \varphi_n(X) - \varphi_n(S_0).$$

Since $X \setminus S_0 = \bigsqcup_{S_i \neq S_0} S_i$ is again a stratification, by induction on the number of strata we have

$$\varphi_n(X) = \sum \varphi_n(S_i).$$

Let X be an arbitrary singular variety with $\dim X \leq n$. Consider two stratifications $X = \bigsqcup S_i$, $X = \bigsqcup S'_j$ of X . Suppose the second stratification refines the first; that is, each stratum S_i is a union of strata S'_j . Then by the additivity of φ_n for nonsingular varieties, we have $\sum \varphi_n(S_i) = \sum \varphi_n(S'_j)$. In general, given two stratification of X there exists a third stratification refining both of them. This shows that the definition (ii) of $\varphi_n(X)$ is independent of the stratification.

We now show the additivity (c) of φ_n . Suppose $\dim X \leq n$, and let Y be a subvariety of X . There exists a stratification of X such that Y is a union of strata. So the additivity of φ_n follows from the additivity of φ_n for nonsingular varieties. This completes the inductive proof of (F_n) .

Clearly the φ_n defined above, $n \geq 0$, give a unique extension of e to a group homomorphism $\varphi : K_0(\mathcal{V}_{\mathbb{R}}) \rightarrow G$. If e is a ring homomorphism then it is easy to see from (i) and (ii) above, by induction on dimension, that φ is also a ring homomorphism. $\square$

2. THE VIRTUAL POINCARÉ POLYNOMIAL

Let X be a nonsingular compact real algebraic variety, and for $i \geq 0$ let $b_i(X)$ be the i th Betti number with $\mathbb{Z}_2$ coefficients,

$$b_i(X) = \dim_{\mathbb{Z}_2} H^i(X; \mathbb{Z}_2) = \dim_{\mathbb{Z}_2} H_i(X; \mathbb{Z}_2).$$

Proposition 2.1. *Let $\pi : \text{Bl}_C X \rightarrow X$ be the blow-up of the nonsingular compact variety X with nonsingular center $C \subset X$. Let E denote the exceptional divisor. Then, for all i ,*

$$b_i(\text{Bl}_C X) - b_i(E) = b_i(X) - b_i(C).$$

Proof. Without loss of generality we may assume that X has pure dimension n and $\dim C < n$.

Let $\tilde{X} = \text{Bl}_C X$. Recall that $H_c^i(X; \mathbb{Z}_2)$ denotes the i th mod 2 cohomology of X with compact supports. Since $\tilde{X} \setminus E \simeq X \setminus C$, the diagram of long exact sequences of closed embeddings

$$\begin{array}{ccccccc} \cdots & \rightarrow & H^{i-1}(E; \mathbb{Z}_2) & \rightarrow & H_c^i(\tilde{X} \setminus E; \mathbb{Z}_2) & \rightarrow & H^i(\tilde{X}; \mathbb{Z}_2) \rightarrow H^i(E; \mathbb{Z}_2) \rightarrow \cdots \\ & & \uparrow \pi^* & & \simeq \uparrow \pi^* & & \uparrow \pi^* & & \uparrow \pi_* \\ \cdots & \rightarrow & H^{i-1}(C; \mathbb{Z}_2) & \rightarrow & H_c^i(X \setminus C; \mathbb{Z}_2) & \rightarrow & H^i(X; \mathbb{Z}_2) \rightarrow H^i(C; \mathbb{Z}_2) \rightarrow \cdots \end{array}$$

induces (by a diagram chase) a long exact sequence

$$(2.1) \quad \cdots \rightarrow H^{i-1}(E; \mathbb{Z}_2) \rightarrow H^i(X; \mathbb{Z}_2) \rightarrow H^i(C; \mathbb{Z}_2) \oplus H^i(\tilde{X}; \mathbb{Z}_2) \rightarrow H^i(E; \mathbb{Z}_2) \rightarrow \cdots$$

Let $\mu_X \in H_n(X; \mathbb{Z}_2)$ denote the fundamental class of X . We have $\pi_*(\mu_{\tilde{X}}) = \mu_X$. Hence by Poincaré duality $\pi^* : H^i(X; \mathbb{Z}_2) \rightarrow H^i(\tilde{X}; \mathbb{Z}_2)$ is injective for all $i \geq 0$. (Let $a \in H^i(X; \mathbb{Z}_2)$. Now $\pi_*(\pi^*a \frown \mu_{\tilde{X}}) = a \frown \pi_*\mu_{\tilde{X}} = a \frown \mu_X$. Thus $\pi^*a = 0$ implies $a \frown \mu_X = 0$, so $a = 0$ by Poincaré duality.) This shows that (2.1) splits into short exact sequences

$$0 \rightarrow H^i(X; \mathbb{Z}_2) \rightarrow H^i(C; \mathbb{Z}_2) \oplus H^i(\tilde{X}; \mathbb{Z}_2) \rightarrow H^i(E; \mathbb{Z}_2) \rightarrow 0,$$

so

$$b_i(C) + b_i(\tilde{X}) = b_i(X) + b_i(E),$$

as desired. $\square$

Corollary 2.2. *For each nonnegative integer i there exists a unique group homomorphism $\beta_i : K_0(\mathcal{V}_{\mathbb{R}}) \rightarrow \mathbb{Z}$ such that $\beta_i(X) = b_i(X)$ for X compact nonsingular.*

There exists a unique ring homomorphism $\beta(\cdot, t) : K_0(\mathcal{V}_{\mathbb{R}}) \rightarrow \mathbb{Z}[t]$ such that $\beta(X, t) = \sum_i b_i(X)t^i$ for X compact nonsingular.

Proof. By the Künneth formula, the classical Poincaré polynomial $P(X, t) = \sum_i b_i(X)t^i$ is multiplicative for compact nonsingular varieties. Thus the corollary follows from Proposition 2.1 and Theorem 1.3. $\square$

Definition 2.3. The integer $\beta_i(X)$ is the i th *virtual Betti number* of the real algebraic variety X , and the polynomial $\beta(X, t) = \sum_i \beta_i(X)t^i$ is the *virtual Poincaré polynomial* of X .

Theorem 2.4. *The virtual Poincaré polynomial $\beta(X, t)$ is of degree $n = \dim X$, and $\beta_n(X) > 0$. In particular, $[X] = [Y]$ implies $\dim X = \dim Y$, and $[X] \neq 0$ if $X \neq \emptyset$.*

Proof. The proof of Theorem 1.3 gives us formulas for $\beta_i(X)$ in terms of a nonsingular compactification of X (1.3)(i), or in terms of a stratification of X (1.3)(ii). Using these formulas we establish the theorem by induction on the dimension of X .

If $\dim X = 0$ then X is a finite set of points, so $\beta(X, t) = a$, where a is the number of points of X .

Suppose the theorem is true for varieties of dimension less than n , and X has dimension n . First suppose X is nonsingular. Let $\bar{X}$ be a nonsingular compactification of X , and let $D = \bar{X} \setminus X$. Then

$$\beta(X, t) = \beta(\bar{X}, t) - \beta(D, t).$$

Since $\dim D < n$, the polynomial $\beta(D, t)$ has degree less than n by inductive hypothesis. Since $\bar{X}$ is compact and nonsingular, $\beta(\bar{X}, t)$ is the Poincaré polynomial of $\bar{X}$, which is of degree n , with $\beta_n(\bar{X}) = b_n(\bar{X}) > 0$. Thus $\beta(X, t)$ has degree n and $\beta_n(X) > 0$.

Now if X is an arbitrary variety of dimension n , let $X = \bigsqcup S_i$ be a stratification of X by nonsingular varieties S_i . Then

$$\beta(X, t) = \sum_i \beta(S_i, t).$$

Since each stratum S_i is nonsingular of dimension at most n , we have that $\beta(S_i, t)$ has degree at most n , and if $\beta(S_i, t)$ has degree n then $\beta_n(S_i) > 0$. So $\beta(X, t)$ has degree n and $\beta_n(X) > 0$. $\square$

Remark 2.5. In [21] Quarez studies the Grothendieck ring $K_0(\mathcal{S})$ of semialgebraic sets, which is generated by homeomorphism classes of semialgebraic sets with sum relation 1.1(2) and product given by 1.1(3). He observes that $[X] = [Y]$ in $K_0(\mathcal{S})$ if and only if $\chi_c(X) = \chi_c(Y)$. In particular, the class of a non-empty semialgebraic set can be zero in $K_0(\mathcal{S})$, and two semialgebraic sets of different dimensions may represent the same class in $K_0(\mathcal{S})$. This makes the construction of motivic measures—more precisely, completion with respect to virtual dimension—impossible for $K_0(\mathcal{S})$.

The generalized Euler characteristics

$$\begin{aligned} \beta(X, -1) &= \sum_i (-1)^i \beta_i(X) \\ \chi_c(X) &= \sum_i (-1)^i \dim H_c^i(X; \mathbb{Z}_2) \end{aligned}$$

are equal for X compact and nonsingular—they both equal the Euler characteristic $\chi(X)$. It follows from Theorem 1.3 that $\beta(X, -1) = \chi_c(X)$ for all real algebraic varieties X . But in general $\beta_i(X) \neq \dim H_c^i(X; \mathbb{Z}_2)$. In fact $\beta_i(X)$ can be negative for $i < \dim X$, and the virtual Betti numbers $\beta_i(X)$ are not topological invariants.

Example 2.6. Consider the union of two intersecting ellipses,

$$X = \{(x, y) \mid (2x^2 + y^2 - 1)(x^2 + 2y^2 - 1) = 0\}.$$

Let X_1 and X_2 be the two irreducible components of X . Then

$$\beta_0(X) = \beta_0(X_1) + \beta_0(X_2) - \beta_0(X_1 \cap X_2) = -2.$$

Example 2.7. For the “figure eight” curve

$$X = \{(x, y) \mid y^2 = x^2 - x^4\},$$

the proper transform of X under the blowup of the plane at the origin is homeomorphic to a circle, and the preimage of the singular point of X is two points. It follows that $\beta_1(X) = 1$. Now consider a second “figure eight” curve

$$Y = \{(x, y) \mid ((x+1)^2 + y^2 - 1)((x-1)^2 + y^2 - 1) = 0\}.$$

Since Y is the union of two circles tangent at the origin, it follows that $\beta_1(Y) = 2$. But Y is homeomorphic to X .

Example 2.8. Let S^n be the unit n -sphere in $\mathbb{R}^{n+1}$. From the inclusion $\mathbb{R}^{n+1} \subset \mathbb{R}^{n+2}$ we have $S^n \subset S^{n+1}$. From the decomposition $S^{n+1} = (S^{n+1} \setminus S^n) \sqcup S^n$ we see that $\beta_n(S^{n+1} \setminus S^n) = -1$. Now $S^{n+1} \setminus S^n$ is homeomorphic to $\mathbb{R}^{n+1} \sqcup \mathbb{R}^{n+1}$, but $\beta_n(\mathbb{R}^{n+1} \sqcup \mathbb{R}^{n+1}) = 0$. For S^{n+1} is the Alexandroff compactification of $\mathbb{R}^{n+1}$ (cf. [4], p. 76), so $\beta_n(\mathbb{R}^{n+1}) = 0$.

3. COMPLEX VARIETIES

If we apply the complex versions of Theorem 1.3 and Proposition 2.1 to the Poincaré polynomial

$$P_{\mathbb{C}}(X, t) = \sum_i \dim_{\mathbb{C}} H^i(X; \mathbb{C}) t^i,$$

we obtain a generalized Euler characteristic of complex algebraic varieties. This invariant is related to Deligne’s mixed Hodge theory in the following way.

For a complex algebraic variety X , the *E-polynomial* (or *Hodge number characteristic*) is given by

$$E(X, u, v) = \sum_{i,p,q} (-1)^i h_{p,q}^i u^p v^q,$$

where $h_{p,q}^i$ is the dimension of the (p, q) -component of the mixed Hodge structure on $H_c^i(X; \mathbb{C})$. The *E-polynomial* is a generalized Euler characteristic (see [8], [12], [20], [7]). Therefore so is the *weight characteristic*

$$(3.1) \quad E(X, t, t) = \sum_{i,j} (-1)^i w_j^i t^j.$$

The coefficients of the weight characteristic are given by

$$w_j^i(X) = \sum_{p+q=j} h_{p,q}^i(X) = \dim_{\mathbb{C}} W_j^i(X) / W_{j-1}^i(X),$$

where

$$(3.2) \quad 0 \subset W_0^i(X) \subset W_1^i(X) \subset \cdots \subset W_i^i(X) = H_c^i(X; \mathbb{C})$$

is the *weight filtration* of cohomology with compact supports [9].

If X is compact and nonsingular then $W_{i-1}^i = 0$, so

$$w_j^i(X) = \begin{cases} b_i^{\mathbb{C}}(X) & i = j \\ 0 & i \neq j, \end{cases}$$

where $b_i^{\mathbb{C}}(X) = \dim_{\mathbb{C}} H^i(X; \mathbb{C})$, and thus the weight characteristic is the Poincaré polynomial $P_{\mathbb{C}}(X, -t)$. In other words, for complex algebraic varieties the virtual Poincaré polynomial is the weight characteristic evaluated at $-t$, and so the j th virtual Betti number is $(-1)^j$ times the *weight j Euler characteristic*,

$$\beta_j^{\mathbb{C}}(X) = (-1)^j \sum_i (-1)^i w_j^i(X).$$

The weight j Euler characteristic has been studied by Durfee [12]. The virtual Poincaré polynomial has been used by Fulton ([13], p. 92) to compute Betti numbers of toric varieties.

Remark 3.1. Applying the complex versions of Theorem 1.3 and Proposition 2.1 to the Poincaré polynomial with coefficients in $\mathbb{Z}_p$, p prime, we get further examples of generalized Euler characteristics of complex algebraic varieties. The existence of these invariants also follows from the work of Gillet and Soulé [14].

Remark 3.2. One could study the Grothendieck ring of real algebraic varieties by means of complexification, that is by considering the isomorphism classes of pairs $(X_{\mathbb{C}}, \tau)$, where $X_{\mathbb{C}}$ is a complex algebraic variety with complex conjugation τ . The real algebraic variety $X_{\mathbb{R}}$ associated to $(X_{\mathbb{C}}, \tau)$ is the fix point set of τ . We do not know how the generalized Euler characteristics of $X_{\mathbb{C}}$ and of $X_{\mathbb{R}}$ are related, except for the observation that

$$\chi_c(X_{\mathbb{C}}) \equiv \chi_c(X_{\mathbb{R}}) \pmod{2}.$$

Note that a given real algebraic variety admits many different realizations as the fixed point set of conjugation on a complex variety.

4. WEIGHT FILTRATION FOR REAL VARIETIES

It is natural to ask whether the virtual Betti numbers of real algebraic varieties are associated to a weight filtration on mod 2 cohomology with compact supports. Totaro [23] has announced that this is the case. The following examples show that a weight filtration for real varieties cannot have properties as strong as those of the weight filtration for complex varieties.

We analyze two examples. Example 4.2, a curve in $\mathbb{R}^3$, shows that a real weight filtration cannot have both the strict naturality property and the resolution of singularities property enjoyed by the complex weight filtration. Example 4.3, a surface in $\mathbb{R}^3$, shows that there is no natural real weight filtration such that the virtual Betti numbers are the weighted Euler characteristics.

Definition 4.1. A *real weight filtration* W assigns to every real algebraic variety X , and to every $i \geq 0$, a filtration of the i th cohomology of X with compact supports and $\mathbb{Z}_2$ coefficients, of the form

$$0 \subset W_0^i(X) \subset W_1^i(X) \subset \cdots \subset W_i^i(X) = H_c^i(X; \mathbb{Z}_2).$$

If W is a real weight filtration, for all i, j we let

$$w_j^i(X) = \dim_{\mathbb{Z}_2} W_j^i(X) / W_{j-1}^i(X).$$

We will consider the following properties of a real weight filtration. The classical complex weight filtration considered in the previous section has all of these properties.

- (1) *Naturality.* If $f : X \rightarrow Y$ is an algebraic morphism, then for all i and j ,

$$f^*W_j^i(Y) \subset W_j^i(X).$$

- (2) *Strict naturality.* If $f : X \rightarrow Y$ is an algebraic morphism, then for all i and j ,

$$f^*W_j^i(Y) = W_j^i(X) \cap \text{Im } f^*.$$

- (3) *Manifold.* If X is compact and nonsingular, then for all i

$$W_{i-1}^i(X) = 0.$$

- (4) *Resolution.* If X is compact and $p : \tilde{X} \rightarrow X$ is a resolution of singularities, then for all i

$$W_{i-1}^i(X) = \text{Ker } p^*.$$

- (5) *Virtual Betti.* For all X and all $i, j \geq 0$, the virtual Betti number β_i is given by

$$\beta_j(X) = (-1)^j \sum_i (-1)^i w_j^i(X).$$

- (6) *Mayer-Vietoris.* If A and B are closed subvarieties of X with $A \cup B = X$ and $A \cap B = C$, then for all j the Mayer-Vietoris cohomology sequence restricts to an exact sequence

$$\cdots \rightarrow W_j^i(X) \rightarrow W_j^i(A) \oplus W_j^i(B) \rightarrow W_j^i(C) \rightarrow W_j^{i+1}(X) \rightarrow \cdots$$

- (7) *Pair.* If A is a closed subvariety of X , then for all j the exact cohomology sequence of the pair restricts to an exact sequence

$$\cdots \rightarrow W_j^i(X \setminus A) \rightarrow W_j^i(X) \rightarrow W_j^i(A) \rightarrow W_j^{i+1}(X \setminus A) \rightarrow \cdots$$

Clearly condition (2) implies (1), (4) implies (3), (7) and (3) imply (5). Since the virtual Betti numbers of a compact nonsingular variety equal the classical Betti numbers, the virtual Betti condition (5) implies the manifold condition (3).

In the rest of this section we will write $H^i(X) = H_c^i(X; \mathbb{Z}_2)$. All the varieties we will consider are compact.

Example 4.2. In $\mathbb{R}^3$ with coordinates (x, y, z) , let X be the intersection of the circular cylinder $x^2 + y^2 = 1$ with the union of the two parabolic cylinders $x = z^2$ and $x = -z^2$,

$$X = \{(x, y, z) \mid x^2 + y^2 = 1, (x - z^2)(x + z^2) = 0\}.$$

Thus X is topologically the union of two circles which are tangent at two points. Let $\tilde{X}$ be the disjoint union of the two irreducible components of X , and let $p : \tilde{X} \rightarrow X$ be the resolution of singularities given by the inclusions of the components. Let C be the unit circle in the (x, y) plane, and let $q : X \rightarrow C$ be the projection. Now $\dim H^1(X) = 3$, and the sequence

$$0 \rightarrow H^1(C) \xrightarrow{q^*} H^1(X) \xrightarrow{p^*} H^1(\tilde{X}) \rightarrow 0$$

is exact. A real weight filtration on $H^1(X)$ has the form

$$0 \subset W_0^1(X) \subset W_1^1(X) = H^1(X).$$

If this filtration has the resolution property, then $W_0^1(X) = \text{Ker } p^* = \text{Im } q^*$. Thus strict naturality of the weight filtration implies that $q^*W_0^1(C) = W_0^1(X)$. But the manifold property implies that $W_0^1(C) = 0$, which is a contradiction.

Therefore a real weight filtration cannot satisfy both strict naturality (2) and resolution (4). This example can also be used to show that a real weight filtration cannot satisfy all three of the following conditions: strict naturality, the manifold condition, and the Mayer-Vietoris condition.

To analyze the next example we first restate the virtual Betti condition (5) for surfaces. Let X be a 2-dimensional (compact) real algebraic variety. Consider a real weight filtration of $H^*(X)$,

$$\begin{aligned} 0 &\subset W_0^0(X) = H^0(X) \\ 0 &\subset W_0^1(X) \subset W_1^1(X) = H^1(X) \\ 0 &\subset W_0^2(X) \subset W_1^2(X) \subset W_2^2(X) = H^2(X). \end{aligned}$$

For $i \geq 0$ let b_i be the i th mod 2 Betti number of X , and let β_i be the i th virtual Betti number of X . The non-negative integers $w_j^i = \dim W_j^i(X)/W_{j-1}^i(X)$ satisfy the equations

$$\begin{aligned} (4.1) \quad &w_0^0 = b_0 \\ &w_0^1 + w_1^1 = b_1 \\ &w_0^2 + w_1^2 + w_2^2 = b_2 \\ &w_0^0 - w_0^1 + w_0^2 = \beta_0 \\ &w_1^1 - w_1^2 = \beta_1 \\ &w_2^2 = \beta_2 \end{aligned}$$

Given $b_0, b_1, b_2, \beta_0, \beta_1, \beta_2$, we wish to determine the possible values of w_j^i which satisfy the equations (4.1). If we arrange the w_j^i in an array

w_2^2		
w_1^1	w_1^2	
w_0^0	w_0^1	w_0^2

then (4.1) says that the sum of the i th diagonal is b_i and the alternating sum of the j th row is β_j (cf. the Mayer-Vietoris spectral sequence computation below).

Example 4.3. Let $X = X_1 \cup X_2 \cup X_3$ be the divisor with normal crossings in 3-space with smooth components X_1, X_2, X_3 defined as follows.

Let X_1 be the sphere of radius $\sqrt{2}$ with center $(1, 0, 0)$,

$$X_1 = \{(x, y, z) \mid (x - 1)^2 + y^2 + z^2 = 2\}.$$

Let X_2 be the sphere of radius $\sqrt{2}$ with center $(-1, 0, 0)$,

$$X_2 = \{(x, y, z) \mid (x + 1)^2 + y^2 + z^2 = 2\}.$$

Then $X_1 \cap X_2$ is the circle of radius 1 in the (y, z) plane with center at the origin.

Let X_3 be the torus of revolution which is a tube of radius $\sqrt{2}/2$ with core the circle of radius 4 in the (y, z) plane with center $(0, 4, 0)$,

$$X_3 = \{(x, y, z) \mid x^2 + (\sqrt{(y-4)^2 + z^2} - 4)^2 = 1/2\}.$$

(X_3 is defined by a polynomial equation of degree 4.)

The torus X_3 intersects each of the spheres X_1 and X_2 transversely along a simple closed curve. The torus X_3 meets the circle $X_1 \cap X_2$ transversely at four points p, q, r, s . The locus Σ of singularities of X is the union of the three loops $X_1 \cap X_2$, $X_1 \cap X_3$, $X_2 \cap X_3$, any two of which meet at the four points p, q, r, s . The natural stratification of X consists of the nonsingular points $X \setminus \Sigma$, the double points $\Sigma \setminus \{p, q, r, s\}$, and the triple points $\{p, q, r, s\}$. The stratum $X \setminus \Sigma$ has 17 connected components, 6 on each sphere and 5 on the torus. All but one of these components is a topological 2-cell. The remaining component is a topological annulus. The stratum $\Sigma \setminus \{p, q, r, s\}$ has 12 components, each of which is a 1-cell. So X has a cell structure with 4 0-cells, 13 1-cells, and 17 2-cells, with Euler characteristic 8.

A computation with Mayer-Vietoris sequences (equivalent to the Mayer-Vietoris spectral sequence computation below) gives that $b_0 = 1$, $b_1 = 1$, and $b_2 = 8$. The first homology group of X is generated by a circle on the torus X_3 which is parallel to the core. The second homology group of X is generated by the boundaries of the 8 bounded connected components of $\mathbb{R}^3 \setminus X$.

Now we compute the virtual Betti numbers of X . For $j = 0, 1, 2$ we have

$$\begin{aligned} \beta_j(X) &= \beta_j(X_1) + \beta_j(X_2) + \beta_j(X_3) \\ &\quad - \beta_j(X_1 \cap X_2) - \beta_j(X_1 \cap X_3) - \beta_j(X_2 \cap X_3) \\ &\quad + \beta_j(X_1 \cap X_2 \cap X_3). \end{aligned}$$

Since the $X_i \cap X_j$ are homeomorphic to circles and $X_1 \cap X_2 \cap X_3$ is four points, we have $\beta_0 = 4$, $\beta_1 = -1$, and $\beta_2 = 3$.

Let W be a real weight filtration on the cohomology of X , and suppose that W is natural and satisfies the virtual Betti condition. Since $b_0 = 1$, we have $w_0^0 = 1$. Since $\beta_2 = 3$ we obtain that $w_2^2 = 3$.

Since $b_1 = 1$, we have $w_0^1 + w_1^1 = 1$. Suppose that $w_0^1 = 1$. Then $H^1(X) = W_0^1(X)$. But the inclusion $f : X_3 \rightarrow X$ induces an injection $f^* : H^1(X) \rightarrow H^1(X_3)$. Since X_3 is compact and nonsingular, it follows from the virtual Betti condition that $W_0^1(X_3) = 0$. This contradicts the naturality statement $f^*W_0^1(X) \subset W_0^1(X_3)$.

The remaining option is that $w_1^1 = 1$, which implies by (4.1) that $w_1^2 = 2$ and $w_0^2 = 3$.

The subvariety $X_1 \cup X_2$ of X has $b_0 = 1$, $b_1 = 0$, $b_2 = 3$, $\beta_0 = 1$, $\beta_1 = -1$, $\beta_2 = 2$. By (4.1) we have $w_0^2(X_1 \cup X_2) = 0$, $w_1^2(X_1 \cup X_2) = 1$, and $w_2^2(X_1 \cup X_2) = 2$.

The subvariety $X_1 \cup X_3$ has $b_0 = 1$, $b_1 = 2$, $b_2 = 3$, $\beta_0 = 1$, $\beta_1 = 1$, $\beta_2 = 2$. Now $w_0^1(X_3) = 0$, $w_1^1(X_3) = 2$, and the restriction $H^1(X_1 \cup X_3) \rightarrow H^1(X_3)$ is an isomorphism. Therefore, by naturality of W , $w_0^1(X_1 \cup X_3) = 0$. By (4.1) we conclude that $w_0^2(X_1 \cup X_3) = 0$, $w_1^2(X_1 \cup X_3) = 1$, and $w_2^2(X_1 \cup X_3) = 2$. The same computation applies to $X_2 \cup X_3$.

There exists a basis $\{a_1, a_2, a_3, a_{12}, a_{13}, a_{23}, b, c\}$ of $H^2(X)$ such that each a_i restricts to the generator of $H^2(X_i)$, and each a_{ij} restricts to a generator of $W_1^2(X_i \cup X_j)$. (The class dual to a_i in homology is represented by the surface X_i , and the class dual to a_{ij} in homology is

represented by the boundary of the intersection of the region inside X_i and the region inside X_j .) Let $f_i : X_i \rightarrow X$ and $f_{ij} : X_i \cup X_j \rightarrow X$, $1 \leq i < j \leq 3$, be the inclusions. By naturality of W , for $k = 0, 1, 2$,

$$f_i^* W_k^2(X) \subset W_k^2(X_i), \quad f_{ij}^* W_k^2(X) \subset W_k^2(X_i \cup X_j).$$

It follows that the images of a_1, a_2, a_3 form a basis for $W_2^2(X)/W_1^2(X)$, and the images of a_{12}, a_{13}, a_{23} are independent in $W_1^2(X)/W_0^2(X)$. Therefore $w_1^2 \geq 3$, which is a contradiction.

Therefore a real weight filtration cannot satisfy both naturality (1) and the virtual Betti condition (5). This example can also be used to show that a real weight filtration cannot satisfy both the Mayer-Vietoris condition and the virtual Betti condition.

It is instructive to compute for Example 4.3 the Mayer-Vietoris spectral sequence which expresses the cohomology of X in terms of the cohomology of its irreducible components X_1, X_2, X_3 . Let X^0 be the disjoint union of X_1, X_2 , and X_3 . Let X^1 be the disjoint union of $X_1 \cap X_2, X_1 \cap X_3$, and $X_2 \cap X_3$. Let $X^2 = X_1 \cap X_2 \cap X_3$.

The Mayer-Vietoris spectral sequence is the spectral sequence of the double complex $C^q(X^p)$, where C^q is the group of singular cochains (mod 2). It converges to the cohomology of X (cf. [15], ch. 2, §5). We have $E_1^{pq} = H^q(X^p)$. Since each $X_i \cap X_j$ is homeomorphic to a circle and $X_1 \cap X_2 \cap X_3$ is four points, we obtain the E_1 term:

$(\mathbb{Z}_2)^3$		
$(\mathbb{Z}_2)^2$	$(\mathbb{Z}_2)^3$	
$(\mathbb{Z}_2)^3$	$(\mathbb{Z}_2)^3$	$(\mathbb{Z}_2)^4$

Computing the differential $d_1 : E_1^{pq} \rightarrow E_1^{p+1,q}$ we find E_2 :

$(\mathbb{Z}_2)^3$		
$(\mathbb{Z}_2)^2$	$(\mathbb{Z}_2)^3$	
$\mathbb{Z}_2$	0	$(\mathbb{Z}_2)^3$

The only possible non-zero differential $d_2^{0,1} : E_2^{0,1} \rightarrow E_2^{2,0}$ has rank 1, and hence we obtain E_3 :

$(\mathbb{Z}_2)^3$		
$\mathbb{Z}_2$	$(\mathbb{Z}_2)^3$	
$\mathbb{Z}_2$	0	$(\mathbb{Z}_2)^2$

Finally, $d_r = 0$ for $r \geq 3$, so $E_3 = E_\infty$. This computation gives that $b_0 = 1$, $b_1 = 1$, and $b_2 = 8$, as claimed above.

The filtration W of the cohomology of X corresponding to the Mayer-Vietoris spectral sequence has $w_j^i = \dim E_\infty^{i-j,j}$. The alternating sum of the ranks of the entries of the j th row of E_1 is the virtual Betti number β_j . Because the j th row of E_2 is the homology of the j th row of E_1 , the alternating sum of the ranks of the entries of the j th row of E_2 is also β_j . (This holds for the Mayer-Vietoris spectral sequence of any divisor with normal crossings.) Since

the differential d_2 is non-zero, the terms E_r for $r > 2$ no longer have this property. Thus the filtration associated to the Mayer-Vietoris spectral sequence does not satisfy the virtual Betti condition.

REFERENCES

- [1] D. Abramovich, K. Karu, K. Matsuki, J. Włodarczyk, *Torification and factorization of birational maps*, J. Amer. Math. Soc. **29** (2002), 531–572.
- [2] V. Batyrev, *Non-Archimedean integrals and stringy Euler numbers of log-terminal pairs*, J. Eur. Math. Soc. **1** (1999), 5–33.
- [3] F. Bittner, *The universal Euler characteristic for varieties of characteristic zero*, arXiv:math.AG/0111062.
- [4] J. Bochnak, M. Coste, M.-F. Roy, *Real Algebraic Geometry*, Springer-Verlag, Berlin 1998.
- [5] A. Borel, J. C. Moore, *Homology theory for locally compact spaces*, Michigan Math. J. **7** (1960), 137–159.
- [6] A. Borel *et al.*, *Intersection Cohomology*, Progress in Math. **50**, Birkhäuser, Boston 1984.
- [7] A. Craw, *An introduction to motivic integration*, arXiv:math.AG/9911179.
- [8] V. Danilov, A. Khovanskii, *Newton polyhedra and an algorithm for computing Hodge-Deligne numbers*, Math. USSR Izvestiya **29** (1987), 279–298.
- [9] P. Deligne, *Poids dans la cohomologie des variétés algébriques*, Proc. Int. Cong. Math. Vancouver (1974), 79–85.
- [10] J. Denef, F. Loeser, *Germes of arcs on singular algebraic varieties and motivic integration*, Invent. Math. **135** (1999), 201–232.
- [11] J. Denef, F. Loeser, *Geometry of arc spaces of algebraic varieties*, Eur. Cong. Math. **1** (2001), 325–348.
- [12] A. Durfee, *Algebraic varieties which are a disjoint union of subvarieties*, Geometry and Topology: Manifolds, Varieties and Knots, C. McCrory, T. Shifrin (eds.), Marcel Dekker, New York 1987, 99–102.
- [13] W. Fulton, *Introduction to Toric Varieties*, Annals of Math. Studies **131**, Princeton 1993.
- [14] H. Gillet, C. Soulé, *Descent, motives, and K-theory*, J. Reine Angew. Math. **478** (1996), 127–176.
- [15] R. Godement, *Topologie Algébrique et Théorie des Faisceaux*, Hermann, Paris 1964.
- [16] M. Goresky, R. MacPherson, *Intersection homology II*, Invent. Math. **71** (1983), 77–129.
- [17] H. Hironaka, *Triangulations of algebraic sets*, A.M.S. Proc. Symp. Pure Math. **29** (1975), 165–185.
- [18] M. Kontsevich, lecture at Orsay, December 7, 1995.
- [19] S. Łojasiewicz, *Triangulation of semi-analytic sets*, Ann. Sc. Norm. Sup. Pisa **18** (1964), 449–474.
- [20] E. Looijenga, *Motivic measures*, Séminaire Bourbaki, exposé 874, March 2000.
- [21] R. Quarez, *Espace des germes d’arcs réels et série de Poincaré d’un ensemble semi-algébrique*, Ann. Inst. Fourier **51** (2001), 43–67.
- [22] J.-P. Serre, *Faisceaux algébriques cohérents*, Annals of Math. **61** (1955), 197–278.
- [23] B. Totaro, *Topology of singular algebraic varieties*, Proc. Int. Cong. Math. Beijing, vol. 1 (2002), 533–541.

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ D’ANGERS, 2 BD LAVOISIER, 49045 ANGERS CEDEX, FRANCE

E-mail address: `parus@tonton.univ-angers.fr`

MATHEMATICS DEPARTMENT, UNIVERSITY OF GEORGIA, ATHENS GA 30602, USA

E-mail address: `clint@math.uga.edu`